

**GUJARAT TECHNOLOGICAL UNIVERSITY****BE - SEMESTER- IV EXAMINATION – SUMMER 2020****Subject Code: 2140706****Date: 29/10/2020****Subject Name: NUMERICAL AND STATISTICAL METHODS FOR  
COMPUTER ENGINEERING****Time: 10:30 AM TO 01:00 PM****Total Marks: 70****Instructions:**

1. Attempt all questions.
2. Make suitable assumptions wherever necessary.
3. Figures to the right indicate full marks.

- |   | <b>MARKS</b> |     |     |     |    |   |     |     |     |     |   |   |  |
|---|--------------|-----|-----|-----|----|---|-----|-----|-----|-----|---|---|--|
| <b>Q.1 (a)</b> Find the relative error if the number $X = 0.004997$ is  | <b>03</b>    |     |     |     |    |   |     |     |     |     |   |   |  |
| (i) truncated to three decimal places.  |              |     |     |     |    |   |     |     |     |     |   |   |  |
| (ii) rounded off to three decimal places.   |              |     |     |     |    |   |     |     |     |     |   |   |  |
| <b>(b)</b> Find the negative root of $x^3 - 7x + 3 = 0$ by the bisection method correct up to three decimal places.   | <b>04</b>    |     |     |     |    |   |     |     |     |     |   |   |  |
| <b>(c)</b> Using Gauss Jacobi method solve the following system of the equations:   | <b>07</b>    |     |     |     |    |   |     |     |     |     |   |   |  |
| $8x - y + 2z = 13$  |              |     |     |     |    |   |     |     |     |     |   |   |  |
| $x - 10y + 3z = 17$   |              |     |     |     |    |   |     |     |     |     |   |   |  |
| $3x + 2y + 12z = 25$  |              |     |     |     |    |   |     |     |     |     |   |   |  |
| <br><b>Q.2 (a)</b> Using trapezoidal rule to evaluate $\int_0^2 \frac{x}{\sqrt{2+x^2}} dx$ , dividing the interval into four equal parts.   | <b>03</b>    |     |     |     |    |   |     |     |     |     |   |   |  |
| <b>(b)</b> By using Lagrange's interpolation formula, find $y(10)$ .  | <b>04</b>    |     |     |     |    |   |     |     |     |     |   |   |  |
| <table border="1" style="border-collapse: collapse; width: 100%; text-align: center;"> <tr> <td style="width: 15%;">x</td> <td style="width: 15%;">5</td> <td style="width: 15%;">6</td> <td style="width: 15%;">9</td> <td style="width: 15%;">11</td> </tr> <tr> <td>y</td> <td>12</td> <td>13</td> <td>14</td> <td>16</td> </tr> </table>                                      | x            | 5   | 6   | 9   | 11 | y | 12  | 13  | 14  | 16  |   |   |  |
| x   | 5            | 6   | 9   | 11  |    |   |     |     |     |     |   |   |  |
| y   | 12           | 13  | 14  | 16  |    |   |     |     |     |     |   |   |  |
| <b>(c)</b> Using the Runge-Kutta method of fourth order, solve $10 \frac{dy}{dx} = x^2 + y^2$ , $y(0) = 1$ at $x = 0.1, x = 0.2$ taking $h = 0.1$   | <b>07</b>    |     |     |     |    |   |     |     |     |     |   |   |  |
| <b>OR</b>   |              |     |     |     |    |   |     |     |     |     |   |   |  |
| <b>(c)</b> Using Euler's method find the approximate value of $y$ at $x = 1.5$ taking $h = 0.1$ . Given that $\frac{dy}{dx} = \frac{y-x}{\sqrt{xy}}$ and $y(1) = 2$ .   | <b>07</b>    |     |     |     |    |   |     |     |     |     |   |   |  |
| <br><b>Q.3 (a)</b> Using Newton Raphson method find the positive root of $x^4 - x - 10 = 0$ correct up to three decimal places.   | <b>03</b>    |     |     |     |    |   |     |     |     |     |   |   |  |
| <b>(b)</b> Fit a least square quadratic curve to the following data:  | <b>04</b>    |     |     |     |    |   |     |     |     |     |   |   |  |
| <table border="1" style="border-collapse: collapse; width: 100%; text-align: center;"> <tr> <td style="width: 15%;">x</td> <td style="width: 15%;">1</td> <td style="width: 15%;">2</td> <td style="width: 15%;">3</td> <td style="width: 15%;">4</td> </tr> <tr> <td>y</td> <td>1.7</td> <td>1.8</td> <td>2.3</td> <td>3.2</td> </tr> </table> Estimate $y(2.4)$ .               | x            | 1   | 2   | 3   | 4  | y | 1.7 | 1.8 | 2.3 | 3.2 |   |   |  |
| x   | 1            | 2   | 3   | 4   |    |   |     |     |     |     |   |   |  |
| y   | 1.7          | 1.8 | 2.3 | 3.2 |    |   |     |     |     |     |   |   |  |
| <b>(c)</b> Find the regression coefficients $b_{yx}$ and $b_{xy}$ hence, find the correlation coefficient between $x$ and $y$ for the following data  | <b>07</b>    |     |     |     |    |   |     |     |     |     |   |   |  |
| <table border="1" style="border-collapse: collapse; width: 100%; text-align: center;"> <tr> <td style="width: 15%;">x</td> <td style="width: 15%;">4</td> <td style="width: 15%;">2</td> <td style="width: 15%;">3</td> <td style="width: 15%;">4</td> <td style="width: 15%;">2</td> </tr> <tr> <td>y</td> <td>2</td> <td>3</td> <td>2</td> <td>4</td> <td>4</td> </tr> </table> | x            | 4   | 2   | 3   | 4  | 2 | y   | 2   | 3   | 2   | 4 | 4 |  |
| x   | 4            | 2   | 3   | 4   | 2  |   |     |     |     |     |   |   |  |
| y   | 2            | 3   | 2   | 4   | 4  |   |     |     |     |     |   |   |  |

OR

- Q.3** (a) Using Simpson's 1/3 rule, find  $\int_0^{0.6} e^{-x^2} dx$ , by taking  $n = 6$ . **03**
- (b) Using Newton's divided difference formula, compute  $f(10.5)$  from the following data: **04**

x	10	11	13	17
f(x)	2.3026	2.3979	2.5649	2.8332

- (c) Solve  $x^4 - 8x^3 + 39x^2 - 62x + 50$  by using Lin Bairstow method up to third iteration starting with  $p_0 = q_0 = 0$ . **07**
- Q.4** (a) Find a real root of the equation  $x \log_{10} x = 1.2$  by the regula falsi method. **03**
- (b) The first four moments of distribution about  $x = 2$  are 1, 2.5, 5.5 and 16. Calculate the four moments about  $\bar{x}$  and about zero. **04**
- (c) Given that  $2 \frac{dy}{dx} = y^2 + x^2 y^2$ ,  $y(0) = 1$ ,  $y(0.1) = 1.06$ ,  $y(0.2) = 1.12$ ,  $y(0.3) = 1.21$  evaluate  $y(0.4)$  by Milne's predictor-corrector method. **07**

OR

- Q.4** (a) Find the arithmetic mean from the following data: **03**

Marks less than	10	20	30	40	50	60
No. of students	10	30	60	110	150	180

- (b) (i) Obtain relation between  $\Delta$  and E. **04**  
(ii) Obtain relation between D and E.
- (c) Obtain cubic spline for every subinterval from the following data **07**

x	0	1	2	3
f(x)	1	2	33	244

- Q.5** (a) Two unbiased coins are tossed. Find expected value of number of heads. **03**
- (b) By Simpson's 3/8 rule, evaluate  $\int_0^1 \frac{\sin x}{x} dx$  taking  $h = \frac{1}{6}$ . **04**
- (c) From the following table, estimate the number of students who obtained marks between 40 and 45. **07**

Marks	30-40	40-50	50-60	60-70	70-80
No. of students	31	42	51	35	31

OR

- Q.5** (a) Using Budan's theorem find the number of roots of the equation  $f(x) = x^4 - 4x^3 + 3x^2 - 10x + 8 = 0$  in the interval  $[-1, 0]$ . **03**
- (b) Find the positive solution of  $x - 2 \sin x = 0$ , correct up to three decimal places starting from  $x_0 = 2$  and  $x_1 = 1.9$ . Using secant method. **04**
- (c) Using Gauss Siedel method solve the following system of the equations: **07**
- $$3x - 0.1y - 0.2z = 7.85$$
- $$0.1x + 7y - 0.3z = -19.3$$
- $$0.3x - 0.2y + 10z = 71.4$$

\*\*\*\*\*